COMPENDIUM OF INTERNATIONAL METHODS OF WINE AND MUST ANALYSIS Maximum acceptable

OIV-MA-C1-01 Maximum acceptable limits of various substances caontained in wine (2019 issue)

Citric acid:	1 g/L	
Volatile acidity:	20 milliequivalents/L The volatile acidity of various specially fortified old wines (wines subject to special legislation and controlled by the government) may exceed this limit.	
Arsenic:	0.2 mg/L	
Boron:	80 mg/L (expressed as boric acid)	
Bromine:	1 mg/L (limit exceeded by way of exception in wines from certain vineyards with a brackish subsoil).	
Cadmium:	0.01 mg/L	
Copper: (OIV-OENO 434-2011)	1 mg/L 2 mg/L for liqueur wines produced from unfermented or slightly fermented grape must	
Diethylene glycol:	п 10 mg/L, to the quantification limit	
Malvidol diglucoside:	15 mg/L (determined by the quantitative method diglucoside described in the Compendium)	
Silver	< 0.1 mg/L	

OIV-MA-C1-01 1

COMPENDIUM OF INTERNATIONAL METHODS OF WINE AND MUST ANALYSIS Maximum acceptable

Total sulfur dioxide at the time of sale to the consumer: (oeno 9/98)				
	 150 mg/L for red wines containing a maximum of 4 g/L of reducing substances. 200 mg/L for white and rosé wines containing a maximum of 4 g/L of reducing substances. 300 mg/L: red, rosé and white wines containing more than 4 g/L of reducing substances. 400 mg/L: in exceptional cases some sweet white wines. 			
Ethanediol /Ethylene glycol:	п 10 mg/L			
Fluoride: (OENO 8/91)	1 mg/L except for wines coming from vineyards treated in conformity with national law, with cryolite in which case, the level of fluoride must not exceed 3 mg/L.			
Methanol: (OENO 19/2004)	400 mg/L for red wines 250 mg/L for white and rosé wines			
Ochratoxin A :	2 μg/L (for wines obtained as from the 2005 harvest)			
(CST 1/2002)				
I I (OW ODNO	0.40 /I for the second and starting from the 2010			
Lead: (OIV-OENO 638-2019)	0.10 mg/L for wine, produced starting from the 2019 harvest year			
,	0.15 mg/L for liqueur wines, produced starting from the 2019 harvest year[1]			
Propan-1,2-diol/propylene glycol (OENO 20/2003)	Still wines : = 150 mg/L Sparkling wines : = 300 mg/L			

OIV-MA-C1-01 2

COMPENDIUM OF INTERNATIONAL METHODS OF WINE AND MUST ANALYSIS Maximum acceptable

Sulfates: (expressed as potassium sulfate)	1 g/L However this limit is raised to:	 П	1.5 g/L
(enpressed as potassiam samate)	- for wines which have undergone a		
	maturing period in casks for at least		
	2 years		
	- for sweetened wines		
	- for wines obtained by the addition		
	to the musts or wine of alcohol or		
	potable spirit		
	for wines with addedconcentrated mustsfor naturally sweet wines	0 0	2.0 g/L
	- for wines obtained under a film "sous voile"	0 0	2.5 g/L

Zinc	5 mg/L	
------	--------	--

^[1] For liqueur wines subject to extended periods of aging, the applicable limit is that in force at the time of their production.

OIV-MA-C1-01 3